

## Footnotes and Financial Disclosures

Originally received: July 17, 2015.

Final revision: November 2, 2015.

Accepted: November 2, 2015.

Available online: December 17, 2015. Manuscript no. 2015-1197.

<sup>1</sup> Singapore Eye Research Institute and Singapore National Eye Centre, Singapore, Republic of Singapore.

<sup>2</sup> Department of Ophthalmology, National University Hospital, National University Health System, Singapore, Republic of Singapore.

Presented as a poster at: European Glaucoma Society Congress, June 2014, Nice, France.

Financial Disclosure(s):

The author(s) have no proprietary or commercial interest in any materials discussed in this article.

Supported by the National Medical Research Council, Singapore, Republic of Singapore (grant no.: NMRC/TCR/002-SERI/2008); and AMO Singapore Pte. Ltd, Singapore, Republic of Singapore (unrestricted grant).

Author Contributions:

Conception and design: Narayanaswamy, Baskaran, Nongpiur, Aung

Analysis and interpretation: Narayanaswamy, Baskaran, Perera, Nongpiur, Htoon, Ho, Aung

Data collection: Narayanaswamy, Baskaran, Perera, Nongpiur, Tun, Wong, Goh, Su, Chew, Ho, Aung

Obtained funding: none

Overall responsibility: Narayanaswamy, Baskaran, Perera, Nongpiur, Htoon, Tun, Wong, Goh, Su, Chew, Aung


Abbreviations and Acronyms:

**ALPI** = argon laser peripheral iridoplasty; **AS OCT** = anterior segment optical coherence tomography; **CI** = confidence interval; **IOP** = intraocular pressure; **LI** = laser iridotomy; **PAC** = primary angle closure; **PACG** = primary angle-closure glaucoma; **PAS** = peripheral anterior synechiae; **PGA** = prostaglandin analog.

Correspondence:

Tin Aung, PhD, FRCOphth, Singapore National Eye Centre, 11 Third Hospital Avenue, Singapore 168751, Republic of Singapore. E-mail: [aung.tin@snecc.com.sg](mailto:aung.tin@snecc.com.sg).

## Pictures & Perspectives



### Bitot's Spots in Gastric Bypass Patient

A 35-year-old female with a history of multiple small bowel resections, presented with several atypical conjunctival neoplasms. She was previously diagnosed with vitamin B<sub>12</sub> deficiency, as well as iron deficiency anemia. A lesion in her right eye demonstrated leukoplakia with a xerotic and foamy appearance. This led to a clinical suspicion of associated vitamin A deficiency (Fig 1). Histologic examination (Fig 2) revealed a keratinized conjunctiva with parakeratosis, hyperkeratosis, and acanthosis. Goblet cells were absent in the area of hyperkeratosis. Histopathologic diagnosis of Bitot's spots was made, correlating her pathological findings with clinical examination.

KRISHNA R. SURAPANENI, MD<sup>1</sup>

AALEYA KOREISHI, MD<sup>2</sup>

PAUL O. PHELPS, MD<sup>1</sup>

<sup>1</sup>University of Wisconsin – Madison, Department of Ophthalmology and Visual Sciences, Madison, Wisconsin; <sup>2</sup>Cornea Consultants of Texas, Fort Worth/Arlington, Texas