

Author Contributions:

Conception and design: Pasquale, Hyman, Danias, Kang

Data collection: Pasquale, Kang

Analysis and interpretation: Pasquale, Hyman, Rosner, Joshipura, Danias, Kang

Obtained funding: Not applicable

Overall responsibility: Pasquale, Hyman, Wiggs, Rosner, Joshipura, McEvoy, McPherson, Danias, Kang

Abbreviations and Acronyms:


**CI** = confidence interval; **HPFS** = Health Professionals Follow-up Study; **HTG** = high-tension glaucoma; **IOP** = intraocular pressure; **MVRR** = multivariable relative risk; **NTG** = normal-tension glaucoma; **POAG** = primary open-angle glaucoma; **VF** = visual field.

Correspondence:

Jae H. Kang, ScD, Channing Division of Network Medicine, 181 Longwood Avenue, Boston, MA 02115. E-mail: [njhjk@channing.harvard.edu](mailto:njhjk@channing.harvard.edu).

## Pictures & Perspectives

---



### Retinal Macroaneurysm in Long-Standing Hypertension

A 74-year-old woman presented with a 6-day history of blurred vision in the left eye. Her vision was 20/20 in the right eye and 20/500 in the left eye. Fundus photograph (Fig 1A) shows superficial and deep retinal hemorrhages associated with a retinal macroaneurysm. Optical coherence tomography angiography (OCTA) demonstrates an outpouching of a retinal arteriole along the inferior arcade (Fig 1B, arrow). Infrared image reveals the structural components of the outpouching and B-scan location (Fig 1C, dotted line). Optical coherence tomography (OCT) at presentation shows the retinal macroaneurysm (Fig 1D, arrow) associated with subretinal and intraretinal fluid. This finding was highly suggestive of long-standing hypertension.

PETER L. NESPER, BA  
EVICA SIMJANOSKI, CRA  
RUKHSANA G. MIRZA, MD

*Department of Ophthalmology, Northwestern University,  
Feinberg School of Medicine, Chicago, Illinois*