

- (IOBA-NHC) in response to hyperosmotic stress. *J Proteome Res.* 2015;14:3982-3995.
30. Vehof J, Zavos HM, Lachance G, et al. Shared genetic factors underlie chronic pain syndromes. *Pain.* 2014;155:1562-1568.

31. Livshits G, Macgregor AJ, Gieger C, et al. An omics investigation into chronic widespread musculoskeletal pain reveals epiandrosterone sulfate as a potential biomarker. *Pain.* 2015;156:1845-1851.

Footnotes and Financial Disclosures

Originally received: August 30, 2016.

Final revision: November 28, 2016.

Accepted: December 8, 2016.

Available online: January 28, 2017. Manuscript no. 2016-179.

¹ Department of Twin Research & Genetic Epidemiology, King's College London, St. Thomas' Hospital, London, United Kingdom.

² Department of Ophthalmology, King's College London, St. Thomas' Hospital, London, United Kingdom.

³ Department of Ophthalmology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

⁴ Department of Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.

Financial Disclosure(s):

The author(s) have made the following disclosure(s): C.J.H.: Grants – Alcon.

Pfizer Worldwide Research and Development, Cambridge, Massachusetts (grant number not applicable) funded the metabolomics screen through an unrestricted grant. The funding organization had no role in the design or conduct of this research. TwinsUK is funded by the Wellcome Trust and the European Community's Seventh Framework Programme (FP7/2007-2013). The study also receives support from the National Institute for Health Research Clinical Research Facility at Guy's and St. Thomas' National Health Service Foundation Trust and National Institute for Health Research

Biomedical Research Centre at Guy's and St. Thomas' National Health Service Foundation Trust and King's College London.

Preliminary data of this study were presented in a paper presentation at: the Association for Research in Vision and Ophthalmology conference, Orlando, Florida, May 6, 2014.

Author Contributions:

Conception and design: Vehof, Hysi, Hammond

Data collection: Vehof, Hysi, Hammond

Analysis and interpretation: Vehof, Hysi, Hammond

Obtained funding: none

Overall responsibility: Vehof, Hysi, Hammond


Abbreviations and Acronyms:

BMI = body mass index; **DED** = dry eye disease; **DHEA** = dehydroepiandrosterone; **DHEA-S** = dehydroepiandrosterone sulfate; **SD** = standard deviation; **SQDES** = Short Questionnaire for Dry Eye Syndrome.

Correspondence:

Jelle Vehof, MD, PhD, Department of Twin Research and Genetic Epidemiology, King's College London, St. Thomas' Hospital Campus, 3rd Floor South Wing Block D, Westminster Bridge Road, London SE1 7EH, United Kingdom. E-mail: jelle.vehof@kcl.ac.uk.

Pictures & Perspectives



Paracentral Acute Middle Maculopathy After Cardiac Arrest

A 33-year-old man suffered a cardiac arrest from ventricular fibrillation complicated by prolonged hypoxia and subsequent ischemia to his brain and liver. After regaining consciousness, he noted blurred vision in his left eye. A diagnosis of paracentral acute middle maculopathy (PAMM) was made. Initially described (Sarraf D, Rahimy E, Fawzi AA, et al. Paracentral acute middle maculopathy: a new variant of acute macular neuroretinopathy associated with retinal capillary ischemia. *JAMA Ophthalmol.* 2013;131:1275-1287) as a hyperreflective band-like lesion at the level of the inner nuclear layer (Fig 1, *dashed arrows*), corresponding to intraretinal whitening visible on funduscopy (*solid arrows*), lesions in PAMM are the result of ischemia to the deep and intermediate capillary plexi that supply the outer levels of the inner retina.

MICHAEL N. COHEN, MD^{1,2}

SABIN DANG, MD^{1,2}

MICHELLE C. LIANG, MD²

¹Ophthalmic Consultants of Boston, Boston, Massachusetts; ²New England Eye Center, Tufts University School of Medicine, Boston, Massachusetts