

- server reproducibility with the portable iCare rebound tonometer and Goldmann applanation tonometer in glaucoma patients. *J Glaucoma*. 2013;22:325-329.
25. Brusini P, Salvétat ML, Zeppieri M, et al. Comparison of iCare tonometer with Goldmann applanation tonometer in glaucoma patients. *J Glaucoma*. 2006;15:213-217.
 26. Sahin A, Basmak H, Niyaz L, Yildirim N. Reproducibility and tolerability of the iCare rebound tonometer in school children. *J Glaucoma*. 2007;16:185-188.
 27. Jablonski KS, Rosentreter A, Gaki S, et al. Clinical use of a new position-independent rebound tonometer. *J Glaucoma*. 2013;22:763-767.
 28. Rao A, Kumar M, Prakash B, Varshney G. Relationship of central corneal thickness and intraocular pressure by iCare rebound tonometer. *J Glaucoma*. 2014;23:380-384.
 29. Munkwitz S, Elkarmouty A, Hoffmann EM, et al. Comparison of the iCare rebound tonometer and the Goldmann applanation tonometer over a wide IOP range. *Graefes Arch Clin Exp Ophthalmol*. 2008;246:875-879.
 30. Calladine D, Packard R. Clear corneal incision architecture in the immediate postoperative period evaluated using optical coherence tomography. *J Cataract Refract Surg*. 2007;33:1429-1435.
 31. Chylack LT, Wolfe JK, Singer DM, et al. The Lens Opacities Classification System III: the Longitudinal Study of Cataract Study Group. *Arch Ophthalmol*. 1993;111:831-836.
 32. Oshika T, Araie M. Time course of changes in aqueous protein concentration and flaw rate after oral acetazolamide. *Invest Ophthalmol Vis Sci*. 1990;31:527-534.

Footnotes and Financial Disclosures

Originally received: July 26, 2016.

Final revision: December 15, 2016.

Accepted: December 17, 2016.

Available online: January 19, 2017.

Manuscript no. 2016-1557.

Hayashi Eye Hospital, Fukuoka, Japan.

Financial Disclosure(s):

The author(s) have made the following disclosure(s): K.H.: Financial support — Alcon Japan Ltd, Santen Pharmaceutical Inc., Wakamoto Pharmaceutical Ltd, Bayer Yakuhin Ltd.

M.Y.: Financial support — Alcon Japan Ltd, Santen Pharmaceutical Inc., Wakamoto Pharmaceutical Ltd, Bayer Yakuhin Ltd.

S.M.: Financial support — Alcon Japan Ltd, Santen Pharmaceutical Inc., Wakamoto Pharmaceutical Ltd, Bayer Yakuhin Ltd.

K.Y.: Financial support — Alcon Japan Ltd, Santen Pharmaceutical Inc., Wakamoto Pharmaceutical Ltd, Bayer Yakuhin Ltd.

Author Contributions:

Conception and design: Hayashi

Analysis and interpretation: Hayashi, Yoshida, Manabe, Yoshimura

Data collection: Yoshida, Manabe, Yoshimura

Obtained funding: none

Overall responsibility: Hayashi


Abbreviations and Acronyms:

AS = anterior segment; IOP = intraocular pressure; OCT = optical coherence tomography; POAG = primary open-angle glaucoma.

Correspondence:

Ken Hayashi, MD, Hayashi Eye Hospital, 4-7-13 Hakataekimae, Hakata-Ku, Fukuoka 812-0011, Japan. E-mail: hayashi-ken@hayashi.or.jp.

Pictures & Perspectives



Unusual Presentation of Choroidal Tuberculoma

A 19-year-old woman with decreased vision in the right eye for 2 months was referred for retinoblastoma. Best corrected visual acuity (BCVA) was 1.9 logMAR. A nodular subretinal lesion with exudative retinal detachment was noted (Fig 1A). Tuberculosis was suspected. Mantoux was 12×12 mm and Quantiferon was positive. Chest computed tomography showed hydro-pyo-pneumothorax with pleural thickening, a partially collapsed left lower lobe, and nodular consolidation with cavitation. Pleural fluid was positive for acid fast bacilli. Head magnetic resonance imaging showed left parietal tuberculoma. Inset: Ultrasonography showing mass lesion with high surface and low internal reflectivity, no calcification. Nine months after systemic therapy, the fundus showed healed tuberculoma with scarring (Fig 1B). BCVA was 0.5 LogMAR.

MAMTA AGARWAL, DNB¹

ADITYA MAITRAY, MS²

VIKAS KHETAN, DO, DNB²

¹Department of Uvea, Sankara Nethralaya, Chennai, Tamil Nadu, India; ²Shri Bhagwan Mahavir Vitreoretinal Services, Sankara Nethralaya, Chennai, Tamil Nadu, India