

Footnotes and Financial Disclosures

Originally received: September 20, 2016.

Final revision: February 16, 2017.

Accepted: February 17, 2017.

Available online: April 3, 2017.

Manuscript no. 2016-379.

¹ Department of Ophthalmology, College of Medicine, University of Ulsan, Asan Medical Center, Seoul, South Korea.

² Department of Applied Statistics, Gachon University, Seongnam-Si, Gyeonggi-do, South Korea.

Financial Disclosure(s):

The author(s) have no proprietary or commercial interest in any materials discussed in this article.

Supported by a grant (2016-0411) from the Asan Institute for Life Sciences, Asan Medical Center, Seoul, South Korea, and by the Basic Science Research Program through the National Research Foundation of Korea, which is funded by the Ministry of Education, Science, and Technology (grant no. NRF-2014R1A1A3A04051089). The funding organization had no part in the study.

Author Contributions:

Conception and design: Kwon, Sung, Han

Data collection: Kwon, Sung

Analysis and interpretation: Kwon, Sung, Han, Moon, Shin

Obtained funding: Not applicable

Overall responsibility: Kwon, Sung, Han


Abbreviations and Acronyms:

AS = anterior segment; **AS-OCT** = anterior segment optical coherence tomography; **CB** = ciliary body; **IOP** = intraocular pressure; **IT₇₅₀** = iris thickness 750 μm from the scleral spur; **LPI** = laser peripheral iridotomy; **LV** = lens vault; **PAC** = primary angle closure; **PACG** = primary angle-closure glaucoma; **PB** = pupillary block; **SS** = scleral spur; **TCA** = trabecular-ciliary process angle; **TCPD** = trabecular-ciliary process distance; **UBM** = ultrasound biomicroscopy; **VF** = visual field.

Correspondence:

Kyung Rim Sung, MD, PhD, Department of Ophthalmology, University of Ulsan, College of Medicine, Asan Medical Center, 388-1 Pungnap-2-dong, Songpa-gu, Seoul, South Korea 138-736. E-mail: sungeye@gmail.com.

Pictures & Perspectives



Purtscher's Retinopathy

A 22-year-old man was ejected from his vehicle sustaining traumatic brain injury, facial fractures, pulmonary contusions, liver laceration, and adrenal hemorrhage. Best-corrected visual acuity was 20/40 in the right eye and hand motions in the left eye. Fundoscopy and color photographs showed numerous cloud-like cotton wool spots surrounding the optic disc bilaterally, and classic intraretinal whitening with clear zones adjacent to the retinal vasculature as seen in the macula of the left eye. Purtscher's retinopathy is an occlusive microangiopathy associated with severe head trauma.

SUNDEEP K. KASI, MD¹

JAMES LARSON, COT²

JAY M. STEWART, MD²

¹The Retina Service of Wills Eye Hospital and Mid Atlantic Retina, Thomas Jefferson University, Philadelphia, Pennsylvania;

²Department of Ophthalmology, University of California San Francisco, San Francisco, California