

Footnotes and Financial Disclosures

Originally received: December 5, 2016.

Final revision: March 20, 2017.

Accepted: March 20, 2017.

Available online: April 19, 2017.

Manuscript no. 2016-959.

Department of Ophthalmology, Seoul National University College of Medicine, Seoul National University Bundang Hospital, Seongnam, Korea.

Financial Disclosure(s):

The author(s) have no proprietary or commercial interest in any materials discussed in this article.

Supported by supported by the Seoul National University Bundang Hospital Research Fund, Seoul, Korea (grant no.: 02-2016-023). The funder had no role in the design or conduct of this research.

Author Contributions:

Conception and design: E.J.Lee, T.-W.Kim

Analysis and interpretation: E.J.Lee, K.M.Lee, S.H.Lee, T.-W.Kim

Data collection: E.J.Lee, K.M.Lee, S.H.Lee, T.-W.Kim

Obtained funding: none

Overall responsibility: E.J.Lee, K.M.Lee, S.H.Lee, T.-W.Kim


Abbreviations and Acronyms:

BP = blood pressure; **D** = diopter; **ICGA** = indocyanine green angiography; **IOP** = intraocular pressure; **ICGPD** = perfusion defect shown by ICGA; **MvD** = microvasculature dropout shown by OCTA; **OCTA** = optical coherence tomography angiography; **POAG** = primary open-angle glaucoma; **RNFL** = retinal nerve fiber layer; **VF** = visual field; **βPPA** = β-parapapillary atrophy.

Correspondence:

Tae-Woo Kim, MD, Department of Ophthalmology, Seoul National University Bundang Hospital, Seoul National University College of Medicine, 82, Gumi-ro, 173 Beon-gil, Bundang-gu, Seongnam, Gyeonggi-do 463-707, Korea. E-mail: twkim7@snu.ac.kr.

Pictures & Perspectives



Tidal Wave Retinal Detachment

A 64-year-old man presented with a 4-week history of progressive vision loss in the right eye. He denied any trauma. His visual acuity was hand motion in the right eye and 20/20 in the left eye. The wide-field fundus photograph (Fig 1A) demonstrated a 6 clock hour giant retinal tear with half the retina folded on itself, bare choroid superotemporally, and a horseshoe tear at 1:30 o'clock. The vertical spectral-domain optical coherence tomography (SD OCT) showed detached macula with the foveal center folded over the inferior retina. The patient underwent pars plana vitrectomy and silicone oil injection. At the 6-week postoperative visit, visual acuity was 20/400 with pinhole. The postoperative fundus photograph (Fig 1B) demonstrated attached retina. The vertical OCT showed silicone oil reflection and ellipsoid zone (arrow) atrophy (between arrowheads) in the foveal center.

JEREMY A. LAVINE, MD, PhD*

JAYA B. KUMAR, MD*

ALEKSANDRA V. RACHITSKAYA, MD

Cole Eye Institute, Cleveland Clinic Foundation, Cleveland, Ohio

*These authors contributed equally to this work